



# GILT CITY

## PROMOTIONAL COUPON OFFERS

Brilliant Distinctions® and Gilt City are offering special coupon offers on JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC treatments.\*

\*Coupons valid for any Brilliant Distinctions® treatments and products.



### APRIL OFFER



Pay \$200, get \$400

towards a JUVÉDERM VOLLURE™ XC treatment

- Promotion begins on GiltCity.com on 4/18/2017
- Following purchase on GiltCity.com, consumer will be directed to Brilliant Distinctions®
- Four (4) \$100 coupons will be deposited
- This offer has no expiration date

### JULY OFFER



Pay \$100, get \$200

towards a JUVÉDERM VOLBELLA® XC treatment

- Promotion begins on GiltCity.com on 7/10/2017
- Following purchase on GiltCity.com, consumer will be directed to Brilliant Distinctions®
- Two (2) \$100 coupons will be deposited
- This offer has no expiration date



### HOW THE COUPONS WORK

- Following purchase, consumers will receive an email with promo codes from GiltCity.com and a link to Brilliant Distinctions®. When the consumer arrives at Brilliant Distinctions®, they will be asked to enter the provided promotional code: 1) after registering as a new member or, 2) after logging in to their existing Brilliant Distinctions® account. Once the codes are successfully entered, coupons will automatically be deposited into their account and appear in Rewards History
- Providers can access Gilt City coupons in the Codes to Redeem section of the Provider Portal when entering a treatment
- Members will receive standard Brilliant Distinctions® Points

- Members can use other available Brilliant Distinctions® coupons towards their purchase and still apply these promotional coupons
- Members can choose how many Gilt City promotional coupons to redeem in a treatment or product purchase; all coupons do not need to be used together
- Full coupon value must be used during same in-office visit
- Coupons can be combined with rebates and never expire
- Each individual can purchase the JUVEDERM VOLLURE™ XC offer twice and the JUVEDERM VOLBELLA® XC offer once. The same individual can redeem the offer(s) at BrilliantDistinctions.com

QUESTIONS? CALL OUR CUSTOMER SERVICE LINE AT 1-888-324-2745.

### INDICATIONS AND IMPORTANT SAFETY INFORMATION

#### JUVÉDERM VOLMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC Important Information

##### INDICATIONS

JUVÉDERM VOLMA® XC injectable gel is indicated for deep (subcutaneous and/or supraperiosteal) injection for cheek augmentation to correct age-related volume deficit in the mid-face in adults over the age of 21.

JUVÉDERM VOLLURE™ XC injectable gel is indicated for injection into the mid-to-deep dermis for correction of moderate to severe facial wrinkles and folds (such as nasolabial folds) in adults over the age of 21.

JUVÉDERM VOLBELLA® XC injectable gel is indicated for injection into the lips for lip augmentation and for correction of perioral rhytids in adults over the age of 21.

##### IMPORTANT SAFETY INFORMATION

##### CONTRAINDICATIONS

These products should not be used in patients who have severe allergies, marked by a history of anaphylaxis or history or presence of multiple severe allergies, and should not be used in patients with a history of allergies to Gram-positive bacterial proteins or lidocaine contained in these products.

##### WARNINGS

• Do not inject into blood vessels. Introduction of these products into the vasculature may lead to embolization, occlusion of the vessels, ischemia, or infarction. Take extra care when injecting soft-tissue fillers;

# JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC

## Important Information (cont.)

for example, inject the product slowly and apply the least amount of pressure necessary. Rare, but serious, adverse events associated with the intravascular injection of soft-tissue fillers in the face have been reported and include temporary or permanent vision impairment, blindness, cerebral ischemia or cerebral hemorrhage leading to stroke, skin necrosis, and damage to underlying facial structures. Immediately stop the injection if a patient exhibits any of the following symptoms: changes in vision, signs of a stroke, blanching of the skin, unusual pain during or shortly after the procedure. Patients should receive prompt medical attention and, possibly, evaluation by an appropriate healthcare professional specialist should an intravascular injection occur.

- Product use at specific sites in which an active inflammatory process (skin eruptions such as cysts, pimples, rashes, or hives) or infection is present should be deferred until the underlying process has been controlled

## PRECAUTIONS

- In order to minimize the risk of potential complications, these products should only be used by healthcare professionals who have appropriate training, experience, and knowledge of facial anatomy
- Healthcare professionals are encouraged to discuss the potential risks of soft-tissue injections with their patients prior to treatment and ensure that patients are aware of signs and symptoms of potential complications
- The safety and effectiveness for the treatment of anatomic regions other than the mid-face with JUVÉDERM VOLUMA® XC, facial wrinkles and folds with JUVÉDERM VOLLURE™ XC, and the lips and perioral area with JUVÉDERM VOLBELLA® XC have not been established in controlled clinical studies
- As with all transcutaneous procedures, dermal filler implantation carries a risk of infection. Follow standard precautions associated with injectable materials
- The safety for use during pregnancy, in breastfeeding females, and in patients with known susceptibility to keloid formation, hypertrophic scarring, and pigmentation disorders has not been studied
- The safety for use of JUVÉDERM VOLUMA® XC in patients under 35 or over 65 years, and JUVÉDERM VOLLURE™ XC and JUVÉDERM VOLBELLA® XC in patients under 22 years, has not been established

- Use with caution in patients on immunosuppressive therapy
- Patients who are using products that can prolong bleeding (such as aspirin, nonsteroidal anti-inflammatory drugs, and warfarin) may experience increased bruising or bleeding at treatment sites
- If laser treatment, chemical peel, or any other procedure based on active dermal response is considered after treatment, or if these products are administered before the skin has healed completely, there is a possible risk of an inflammatory reaction at the treatment site
- Patients who experience skin injury near the site of implantation may be at a higher risk for adverse events
- The safety of JUVÉDERM VOLUMA® XC injectable gel for use in patients with very thin skin in the mid-face has not been established
- Patients may experience late onset nodules with use of dermal fillers, including JUVÉDERM VOLUMA® XC
- Patients may experience late onset adverse events with use of dermal fillers

## ADVERSE EVENTS

The most commonly reported side effects for JUVÉDERM® injectable gels were injection-site redness, swelling, pain, tenderness, firmness, lumps/bumps, bruising, discoloration, and itching. For JUVÉDERM VOLBELLA® XC, dryness was also reported. For JUVÉDERM VOLUMA® XC, side effects were predominantly moderate in severity, with duration of 2 to 4 weeks; for JUVÉDERM VOLLURE™ XC, they were mostly mild or moderate in severity, with duration of 14 days or less; and for JUVÉDERM VOLBELLA® XC, they were predominantly mild or moderate, with duration of 30 days or less.

To report an adverse reaction with any product in the JUVÉDERM® Collection, please call Allergan at 1-800-433-8871. Please visit [JuvedermDFU.com](http://JuvedermDFU.com) for more information.

JUVÉDERM VOLUMA® XC, JUVÉDERM VOLLURE™ XC, and JUVÉDERM VOLBELLA® XC injectable gels are available by prescription only.

